

Discovery of *Maritrema obstipum* (Digenea: Microphallidae) from Migratory Birds in Korea

Ok-Sik Chung¹, Woon-Mok Sohn², Jong-Yil Chai³, Min Seo⁴ and Hye-Jung Lee^{4,*}

¹Division of Ecology and Environment, Chungnam Development Institute, Gongju 314-140, Korea; ²Department of Parasitology, Biomedical Center for Brain Korea 21 an Institute of Health Science Gyeongsang National University School of Medicine, Jinju 660-751, Korea; ³Department of Parasitology and Tropical Medicine, Seoul National University College of Medicine, Seoul 110-799, Korea; ⁴Department of Parasitology and Research Center for Mummy, College of Medicine, Dankook University, Cheonan 330-714, Korea

Abstract: Adults of *Maritrema obstipum* (Digenea: Microphallidae) were found in the intestines of 4 species of migratory birds, including the sanderling (*Crocethia alba*), Kentish plover (*Charadrius alexandrinus*), Mongolian plover (*Charadrius mongolus*), and red-necked stint (*Calidris ruficollis*), collected from Yubu Island, Chungcheongnam-do, Korea. The worms were 451 × 265 µm in size, and were easily identifiable as *Maritrema* species by the presence of the cirrus sac, and the ring-like distribution of the vitellaria. More specifically, the ejaculatory duct curved posteromedially, and the 2 parts of vitelline follicles were found to be distinct at the posterior end. The eggs were brown-colored, and 19.8 × 12.3 µm in size. All these findings implicated *M. obstipum* as the pertinent species of the worms. Beside these, adult worms of *Gynaecotyla squatarolae*, *Parvatrema duboisi*, and *Acanthoparyphium* sp. were also discovered. This is the first report establishing migratory birds as the natural definitive hosts for *M. obstipum*.

Key words: *Maritrema obstipum*, sanderling, Kentish plover, Mongolian plover, red-necked stint

Microphallidae Ward, 1901 is a large family of small flukes found in the intestines of birds, that have consumed crustaceans [1]. In South Korea, the following 4 species have been described from crustacean hosts: *Microphalloides japonicus* from *Helice tridens tridens*, *Levinseniella* spp. from *Macrophthalmus japonicus*, *Gynaecotyla squatarolae*, and *Microphallus koreana* from *Macrophthalmus dilatatus* [2-5]. As for cases of human infections by microphallids, there have been only a limited number of reports on *Spelotrema brevicaca* and *G. squatarolae* [6,7]. Last year, a 5th species of Microphallidae was discovered in an Asian shore crab, *Hemigrapsus sanguineus*, caught in Jebu-do, Gyeonggi-do. An experimental infection revealed adult flukes having a symmetric ribbon-like vitellarium, which feature and other overall detailed morphology identified them as a new species of *Maritrema*, subsequently named *Maritrema jebuensis* [8]. This finding established, for the first time, the presence of *Maritrema* sp. in South Korea. The natural definitive host of *Maritrema* sp., however, was not yet known in South Korea. In

the present study, we discovered the adult worms of *Maritrema obstipum* from the intestine of migratory birds, proving them as the natural definitive hosts of *M. obstipum* in Korea.

In October 2010, 25 dead migratory birds, i.e., 2 sanderlings (*Crocethia alba*), 8 Kentish plover (*Charadrius alexandrinus*), 6 mongolian plover (*Charadrius mongolus*), 8 red-necked stint (*Calidris ruficollis*), and 1 terek sandpiper (*trigna cinereus*) were collected from Yubu-do, Songnim-ri, Janghang-eup, Seochon-gun, Chungcheongnam-do, South Korea. All of the specimens were in relatively good condition and showing no signs of decomposition, and they were transported to the laboratory for further studies. There, the birds' intestines were removed, fixed in 2% formalin and examined under stereomicroscopy. The adult worms found therein were fixed in 10% formalin, and then stained with Semichon's acetocarmine, and finally mounted in permount. Ten specimens each were measured, and the results were recorded in micrometer ranges. Species identification was conducted under optical microscopy.

From the 2 sanderlings, 122 flukes were recovered, 113 of which, on the basis of the symmetric ribbon-like vitellarium, were determined to *Maritrema* sp. All 8 Kentish plovers and all 6 Mongolian plovers were infected with the same species. In the red-necked stints, the infection rate was 12.5% (1/8), and in the case of 1 terek sandpiper, there were no adult worms of

•Received 4 July 2011, revised 21 September 2011, accepted 21 September 2011.

*Corresponding author (kardia84@naver.com)

© 2011, Korean Society for Parasitology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

that species.

The followings are the morphological characters of the recovered *Maritrema* worms (Fig. 1). Body was linguiform, 451 (400-495) $\mu\text{m} \times 265$ (260-270) μm . Suckers were of approximately the same size, the oral 39 \times 39 μm and the ventral 36 \times 39 μm . Prepharynx was absent or, if present, very short. Intestinal ceca extended laterally along the anterior edge of the cirrus sac, almost to the lateral body wall. Genital atrium was sinistral to the ventral sucker near its edge. Cirrus sac, tapering to the prostate region, was 154 (126-178) μm long, with an ejaculatory duct curving posteromedially. The vitellarium consisted of a narrow transverse band of small follicles extending laterally on either side to the body wall, and then turning posteriorly to form a ring around the posterior portion of the body. By magnification, the 2 parts of the vitellaria were found to be distinct at the posterior end. Ovary was ovoid, posterodextral to the ventral sucker and anterior to the vitellarium, and was 49 (37-62) $\mu\text{m} \times 52$ (37-62) μm in size. Testes were lateral, and posterior to the anterior part of vitellaria, and indistinguishable. Eggs were brown-colored and 19.8 \times 12.3 μm in dimensions.

The morphology of *M. obstipum* specimens examined in this study differed from that already reported for *M. jebuensis* [9]. First, in *M. jebuensis*, the vitellaria formed a complete ring, but in *M. obstipum*, the 2 parts of the vitellaria, as discovered in the

present study, were distinct. Second, in *M. obstipum*, the sucker ratio (OS/VS) was found to be 1:1, whereas in *M. jebuensis*, the ventral sucker was larger than the oral sucker. Third, in the present *M. obstipum* species, posteromedial curving of the ejaculatory duct was evident, which had been found not the case for *M. jebuensis*. In light of all of these findings, the worms examined in the present study were identified as belonging to *M. obstipum*, not *M. jebuensis*.

Table 1. Trematodes collected from the intestine of migratory birds

Birds	No. of birds infected with <i>Maritrema obstipum</i> /No. examined	Species of trematodes (number)
Terek sandpiper	0/1	<i>Gynaecotyla squatarolae</i> (4)
Sanderling	2/2	<i>M. obstipum</i> (113) <i>Parvatrema duboisi</i> (2) Unknown trematodes (7)
Kentish plover	8/8	<i>M. obstipum</i> (127) <i>G. squatarolae</i> (6) Unknown trematodes (12)
Mongolian plover	6/6	<i>M. obstipum</i> (73) <i>G. squatarolae</i> (16) Unknown trematodes (52) <i>Acanthoparyphium</i> sp. (3)
Red-necked stint	1/8	<i>M. obstipum</i> (3) <i>G. squatarolae</i> (1) Unknown trematodes (19)




Fig. 1. (A) An adult of *Maritrema obstipum* recovered from the sanderling, *Crocethia alba*. Bar = 100 μm . CS, cirrus sac; VS, ventral sucker; OV, ovary; T, testis; VT, vitellarium; IC, intestinal ceca. (B) A line drawing of the Fig. 1, the adult worm of *M. obstipum*. CS, cirrus sac; ED, ejaculatory duct, ventral sucker; OV, ovary; T, testis; VT, vitellarium.

Genera other than *Maritrema* were also recovered from the migratory birds, as summarized in Table 1. Adult *G. squatarolae* worms were found in 4 bird species, *Parvatrema duboisi* specimens were taken from 1 of the sanderlings. In addition, 3 *Acanthoparyphium* sp. worms were recovered from a Mongolian plover, though species identification was impossible due to the poor condition of those worms.

In South Korea, *M. jebuensis* was, as noted above, first discovered in an Asian shore crab caught in Jebu-do, Gyeonggi-do [8]. It was expected that, as migratory birds have already been identified as definitive hosts for *Maritrema* spp. [10], their role would be essential in South Korea as well. In fact, whereas adult *Maritrema* sp. worms were recovered from 4 species of birds examined in the present study, among which were the Mongolian and Kentish plovers, in which no *M. jebuensis* specimens were found. The absence of *M. jebuensis* in these birds might have reflected the geographic distance between Yubu-do, Chungcheongnam-do and Jebu-do, Gyeonggi-do. In any case, a Yubu-do crab survey to establish the life cycle of *M. obstipum* would be necessary. Certainly, considering the discovery of human *G. squatarolae* infection [7], surveys on *Maritrema* spp. infection, among the residents of Yubu-do, would also be needed.

In a previous study, the ruddy turnstone, *Arenaria interpres interpres*, from the seashore of Gunsan-si, Jeollabuk-do was proven to be a definitive host for *G. squatarolae* [11]. In the present study, in fact, adult *G. squatarolae* worms were also recovered from some species of migratory birds, particularly the terek sandpiper, Kentish plover, and Mongolian plover, thereby establishing them as additional definitive hosts for *G. squatarolae*. A survey on metacercariae of *G. squatarolae* revealed their presence in crabs, especially *M. japonicus*, from the western coastal areas of South Korea [12]. Accordingly, it has been suggested that migratory birds are the sources of *G. squatarolae* infection in those areas. In the present study, worms belonging to the genus *Acanthoparyphium* were discovered in the intestines of Mongolian plovers, but as above-noted, species identification was impossible. Further studies on this genus are required, particularly as *Acanthoparyphium tyosenense* has been established as a human-infecting trematode in South Korea [13].

In South Korea, a total of 19 species of intestinal trematodes, including 12 from the family Heterophyidae, had been reported in humans as of 2002 [14]. Since then, an additional 3 species, specifically *Acanthotrema felis*, *G. squatarolae*, and *Haplorchis pumilio*, have been added to that list [7,15]. Even

though migratory birds are known to play a pivotal role in transmission of parasitic diseases, especially those caused by intestinal trematodes, there have been few reports on such birds in South Korea. The oystercatcher, *Haematopus ostralegus*, has been shown to be a natural definitive host of *Gymnophalloides seoi* [16]. Moreover, as noted above, adult *G. squatarolae* worms were discovered in the intestine of a ruddy turnstone [11], and recently, adult *Parvatrema duboisi* and *Parvatrema homoeotectum* worms were recovered from a great knot and Mongolian plovers [17]. More intestinal flukes, however, have yet to be matched with their definitive hosts. Clearly, further investigation into the migratory birds of South Korea is required.

REFERENCES

1. Bray RA, Gibson DI, Jones A. Keys to the Trematode. Vol. 3. London, UK. CAB International and Natural History Museum. 2008, p 451-492.
2. Seo BS, Yoon JS, Lee CW, Lee SH. Studies on *Microphalloides japonicus*. Korean J Parasitol 1964; 2: 41-51.
3. Choi DW, Lee SW, Kim JM, Kim JK. Studies on the larval trematodes from brackish water crabs: studies on a kind of *Levinseniella*. Korean J Parasitol 1965; 3: 31-38.
4. Seo M, Guk SM, Lee SH, Chai JY. *Gynaecotyla squatarolae* (Digenea: Microphallidae) from rats experimentally infected with metacercariae from the shore crab, *Macrophthalmus dilatatus*. Korean J Parasitol 2007; 45: 199-204.
5. Guk SM, Chai JY, Sohn WM, Kim YM, Sim S, Seo M. *Microphallus koreana* n. sp. (Trematoda: Microphallidae) transmitted by a marine crab, *Macrophthalmus dilatatus*. Korean J Parasitol 2008; 46: 165-169.
6. Africa CM, de Leon W, Garcia EY. Visceral complications in intestinal heterophyidiasis of man. Acta Med Philipp Monogr Ser 1940; 1: 1-132.
7. Chung OS, Lee HJ, Kim YM, Sohn WM, Kwak SJ, Seo M. First report of human infection with *Gynaecotyla squatarolae* and first Korean record of *Haplorchis pumilio* in a patient. Parasitol Int 2011; 60: 227-229.
8. Chung OS, Lee HJ, Sohn WM, Lee SH, Park IY, Oh SA, Chai JY, Seo M. Discovery of *Maritrema jebuensis* n. sp. (Digenea: Microphallidae) from the Asian Shore Crab, *Hemigrapsus sanguineus*, in Korea. Korean J Parasitol 2010; 48: 335-338.
9. Etges FJ. Studies on the life histories of *Maritrema obstipum* (Van Cleave and Mueller, 1932) and *Levinseniella amnicolae* n. sp. (Trematoda: Microphallidae). J Parasitol 1953; 39: 643-662.
10. Coill WH. Notes on the genus *Maritrema* Nicoll, 1907 (Trematoda: Microphallidae) with the description of two new species. J Parasitol 1955; 41: 533-537.
11. Seo M, Guk SM, Chai JY. The ruddy turnstone, *Arenaria interpres interpres*, a new definitive host for *Gynaecotyla squatarolae* (Digenea: Microphallidae) from the Asian Shore Crab, *Hemigrapsus sanguineus*, in Korea. Korean J Parasitol 2010; 48: 335-338.

- nea: Microphallidae). *Korean J Parasitol* 2008; 46: 41-43.
12. Lee HJ, Chai JY, Lee JW, Jin H, Min KH, Cho YJ, Seo M. Surveys of *Gynaecotyla squatarolae* and *Microphallus koreana* (Digenea: Microphallidae) metacercariae in two species of estuarine crabs in western coastal areas, Korea. *Korean J Parasitol* 2010; 48: 81-83.
13. Chai JY, Han ET, Park YK, Guk SM, Lee SH. *Acanthoparyphium tyosenense*: The discovery of human infection and identification of its source. *J Parasitol* 2001; 87: 794-800.
14. Chai JY, Lee SH. Food-borne intestinal trematode infections in the Republic of Korea. *Parasitol Int* 2002; 51: 129-154.
15. Cho SH, Cho PY, Lee DM, Kim TS, Kim IS, Hwang EJ, Na BK, Sohn WM. Epidemiological survey on the infection of intestinal flukes in residents of Muan-gun, Jeollanam-do, the Republic of Korea. *Korean J Parasitol* 2010; 48: 133-138.
16. Ryang YS, Yoo JC, Lee SH, Chai JY. The palearctic oystercatcher *Haematopus ostralegus*, a natural definitive host for *Gymnophalloides seoi*. *J Parasitol* 2000; 86: 418-419.
17. Chung OS, Lee HJ, Sohn WM, Park YK, Chai JY, Seo M. Discovery of *Parvatrema duboisi* and *Parvatrema homoeotecnium* (Digenea: Gymnophallidae) from migratory birds in Korea. *Korean J Parasitol* 2010; 48: 271-274.